

# Assessment of the Role of Fine Needle Aspiration Cytology as Diagnostic Criteria in Follicular Patterned Lesion of Thyroid

Arjun Lal Jain<sup>1</sup>, Sachin Jain<sup>2\*</sup>, Jitendra Acharya<sup>3</sup>

<sup>1</sup>Assistant Professor, Department of Pathology,  
Geetanjali Medical College and Hospital, Udaipur, Rajasthan, India.

<sup>2\*</sup>Assistant Professor, Department of Medicine,  
Ananta Institute of Medical Science and Research Center, Udaipur, Rajasthan, India.

<sup>3</sup>Senior Demonstrator, Department of Dentistry, S.P. Medical College, Bikaner, Rajasthan, India.

## ABSTRACT

**Aim:** The study aimed to evaluate the role of Fine Needle Aspiration Cytology (FNAC) in follicular patterned lesions of thyroid.

**Materials and Methods:** A prospective study was done in 30 cases in which FNAC showed follicular patterned lesions. Histopathological correlation was done wherever possible.

**Results:** Out of 30 cases of follicular patterned lesions 17 cases were reported as follicular neoplasm, 12 cases were reported as adenomatous goiter and one case was reported as follicular variant of papillary carcinoma of thyroid in FNAC. Based on the FNAC diagnosis, surgery was done and histopathological correlation was available in nine cases.

**Conclusion:** FNAC is quite accurate in follicular patterned lesions of thyroid with a sensitivity of 70.6%, specificity of 100% and diagnostic accuracy of 80%.

**Keywords:** FNAC, Follicular Patterned Lesions, Thyroid.

## \*Correspondence to:

**Dr. Sachin Jain,**  
Assistant Professor,  
Department of Medicine,  
Ananta Institute of Medical Science and Research Center,  
Udaipur, Rajasthan, India.

## Article History:

**Received:** 14-09-2017, **Revised:** 01-10-2017, **Accepted:** 19-10-2017

## Access this article online

Website: <a href="http://www.ijmrp.com">www.ijmrp.com</a>	Quick Response code 
DOI: 10.21276/ijmrp.2017.3.6.108	

## INTRODUCTION

Thyroid nodules occur in 20% to 70% of the population, which shows an increasing trend in the incidence with age and comprise a significant proportion of cytology specimens. Fine needle aspiration cytology (FNAC) is the most sensitive, accurate and cost effective initial method for the clinical management of patient with thyroid nodules.<sup>1</sup> Majority of the thyroid nodules are benign, with cancer accounting for only 1 % of all the lesions. Hence it is imperative for the treating surgeon to know the biologic nature, as to benign versus malignant lesions and thus plan appropriate treatment.<sup>2</sup> This would in turn avoid unnecessary surgeries. Follicular patterned lesions are a grey zone area in cytology. This category encompasses wide variety of lesions ranging from benign to malignant lesions such as dominant nodule of an adenomatous goiter, follicular neoplastic lesions including adenoma and carcinoma and follicular variant of papillary carcinoma. The major cause of false positive and false negative rates at FNAC has been attributed to follicular patterned lesions. There are several studies published in literature revealing the difficulties posed by the pathologist in the area of follicular patterned lesions of thyroid.<sup>3,4</sup> In the present study the efficacy of FNAC in the diagnosis of the follicular patterned lesions of thyroid was evaluated.

## MATERIALS AND METHODS

A prospective cross sectional study was conducted over duration of three years. 30 cases depicting follicular patterned lesions at FNAC were selected from the files of cytopathology, irrespective of age and sex of the patient. FNAC was done by non-aspiration technique using a 23 gauge needle. 50% slides were fixed in 50% ethanol and 50% were air dried. The fixed slides were stained with Haematoxylin and Eosin, modified Papanicolaou stains. The air dried smears were stained with Giemsa stain. Two experienced Pathologist reported on the smears in double blinded fashion. Only the adequate smears with predominant follicular cell pattern were included in the study.

The lesions were classified into three categories as adenomatous goiter, follicular neoplasm and follicular variant of papillary carcinoma (FVPCT).

## RESULTS

### Cytology Diagnostic Categories

Among 30 cases of follicular patterned lesions, 17 cases were reported as follicular neoplasm, 12 cases were reported as adenomatous goiter and one case was reported as follicular variant of papillary carcinoma of thyroid. (Table 1)

**Cytology - Histopathologic Correlation**

Histopathological correlation was available in nine cases. Table 2 shows cyto-histopathology correlation in the nine cases. Concordant diagnosis was obtained in six cases. There were two cases of false positive wherein a follicular adenoma was misinterpreted as adenomatous goiter. There was one false negative wherein a case of FVPCT was misdiagnosed as follicular neoplasm. Sensitivity, specificity and accuracy of FNAC in the interpretation of follicular patterned lesions were found to be 70.6%, specificity of 100% and diagnostic accuracy of 80%.

**Table 1: Table depicting cytodiagnostic categories in follicular patterned lesions.**

Categories Of Follicular Patterened Lesion	FNAC Diagnosis N (%)
Adenomatous goiter	12%
Follicular neoplasm	17%
Follicular variant of papillary carcinoma	01%
<b>Total</b>	<b>30</b>

**Table 2: Depicting the Cyto histopathologic correlation**

FNAC	Histopathology	
	Concordent Case	Discordent Cases
FN(4)	03	01
FUPC(1)	01	00
AG(4)	02	02
<b>TOTAL</b>	<b>06</b>	<b>03</b>

FNAC- Fine needle aspiration cytology, FN – Follicular neoplasm, FVPCT- Follicular variant of papillary carcinoma of thyroid, AG-Adenomatous goiter.

**DISCUSSION**

FNAC is a sensitive and specific tool in distinguishing neoplastic from non-neoplastic thyroid lesions. Meticulous examination of the pattern along with cellular details and background characteristics would be helpful in accurately diagnosing thyroid lesions.<sup>4</sup> By light microscopy at FNAC a thyroid lesion is designated as follicular when it totally or almost totally (more than 95%) displays a follicular growth pattern or thyroid follicles with central lumen containing variable amounts of colloid. The differential diagnosis of follicular patterned lesion or neoplasm in thyroid FNAC specimens includes hyperplastic/adenomatous goiter, follicular adenoma and carcinoma and follicular variant of papillary thyroid carcinoma.<sup>4,5</sup> They are diagnosed based on cytoarchitectural pattern, nuclear features, cytological atypia, and background colloid material.

Several studies have shown that the major pitfall of FNAC in thyroid has been the follicular patterned lesions.<sup>6-8</sup> Adenomatous/hyperplastic goiter yields cellular smear with follicular cells in micro and macro follicular pattern. The follicular cells are small and are in the background of colloid. At times dominant nodule of an adenomatous goiter poses a diagnostic challenge if the aspirate shows only repetitive microfollicles. This

can lead to the overdiagnosis as follicular neoplasm. However variable pattern comprising of micro and macrofollicles in other areas of the smear/another smears, background of colloid would aid in the diagnosis of a goitrous lesion. In doubtful cases repeat FNAC from other areas would solve the dilemma. Adenomatous goiter on histopathology shows a complete or partial encapsulation and mixture of micro and macro follicles filled with colloid associated with degenerative changes like hemorrhage, fibrosis and calcification, a morphology which can be appreciated at cytology.<sup>9</sup>

Likewise in follicular neoplasm, aspirate from a macrofollicular area would yield abundant colloid and a variable micro and macrofollicular pattern. This would erroneously label a neoplastic lesion as an adenomatous goiter.

Nevertheless, cytological features like increased cellularity with nuclear crowding and overlapping, repetitive microfollicular pattern, scanty or no colloid can be of aid in distinguishing follicular neoplasms from nodular goiters.<sup>10</sup> It has been proposed that the risk of follicular neoplasm is inversely related to size of the follicle and amount of colloid. Smears with abundant colloid and macrofollicles are at low risk of being neoplastic while smears with scant or no colloid is at high risk.<sup>11</sup> Follicular carcinoma however should have the nuclear features like crowding, overlapping, prominent nucleoli, chromatin clumping and parachromatin clearing.

However at FNAC these lesions would fall under the category of follicular neoplasm. Nothing short of capsular and vascular invasion at histopathology would classify them as follicular carcinoma. In the present study two discordant cases were of follicular adenomas which were diagnosed as adenomatous goiter at FNAC. This discordance was due to the presence of normo/macrofollicles and abundant colloid. Similar pitfall in diagnosis was encountered by Clary et al.<sup>12</sup>

The necrotic material in the aspirate might be misinterpreted as thick colloid. FVPCT showed cell rich smears with follicular epithelial cells in predominantly micro follicular pattern. Occasionally papillary fragments may be noted. The classical nuclear features of papillary carcinoma like nuclear grooving, clearing and intranuclear inclusions and powdery chromatin pattern make the diagnosis evident. However at times fixation error may obscure these findings making the diagnosis difficult. In the present study, we observed similar problem, leading to a false negative diagnosis.<sup>13</sup>

Careful observation of the cell pattern along with cytomorphologic details and background material would classify the follicular patterned lesions into appropriate category, as to non-neoplastic or neoplastic. However at times the grey zone lesions poses dilemma, at such time immunohistochemistry would help the pathologist and surgeons plan appropriate management.

**CONCLUSIONS**

Follicular patterned lesions are a grey zone area in FNAC of thyroid lesions. Careful observation of the cell pattern along with cytomorphologic details and background material would classify the follicular patterned lesions into appropriate category. In the present study FNAC was quite accurate in follicular patterned lesions of thyroid with a sensitivity of 70.6%, specificity of 100% and diagnostic accuracy of 80%.

## REFERENCES

1. Faquin WC. Diagnosis and Reporting of Follicular-Patterned Thyroid Lesions by Fine Needle Aspiration. *Head and neck pathol.* 2009;3:82-85.
2. Lingegowda JB, Muddegowda PH, Rajesh KN, Ramkumar KR: Application of pattern analysis in fine needle aspiration of solitary nodule of thyroid. *J Cytol.* 2010;27:1-7.
3. Zubair W Baloch and Virginia A LiVolsi. Our approach to follicular patterned lesions of the thyroid. *J Clin Pathol.* 2007;60:244–250.
4. Baloch ZW, LiVolsi VA. Follicular-Patterned Lesions of the Thyroid. *AmJ ClinPathol* 2002;117:143-150
5. Pandey P, Dixit A, Mahajan NC. Fine-needle aspiration of the thyroid: A cytohistologic correlation with critical evaluation of discordant cases. 2012;9:32-39
6. Orell SR. In: Orell SR, Sterrett GF, Walters MN, Whitakar D, editors. *Manual and atlas of fine needle aspiration cytology.* 4th ed. New Delhi: Churchill-Livingstone; 2005. p. 125-64.
7. Reagan LR, Farkas T, Dehner LP. Fine needle aspiration of the thyroid: a cytohistologic correlation and study of discrepant cases. *Thyroid* 2004;14:35-41.
8. Basu N, Bhattacharya NK, Chowdhury M, Ghosh AK, Guhamallick M, Roy S, Sengupta S. Cytodiagnosis of thyroid lesions-usefulness and pitfalls: A study of 288 cases. *Journal of Cytology* 2008;25:6-9.
9. Bommanahalli BP, Bhat RV, and Rupanarayan R.A cell pattern approach to interpretation of fine needle aspiration cytology of thyroid lesions: A cyto-histomorphological study. *J Cytol:* 2010; 27:127-32.
10. Atkinson B, Ernest CS, Livolsi V. Cytologic diagnosis of follicular tumours of the thyroid. *Diagn Cytopathol* 1986;2:1-5.
11. Jayaram G. *Atlas and Text of Thyroid Cytology.* 1st ed. New Delhi: 2006.53-54.
12. Clary KM, Condel JL, Liu Y, Johnson DR, Grzybicki DM, Raab SS. Interobserver variability in the fine needle aspiration biopsy diagnosis of follicular lesions of the thyroid gland. *Acta Cytol.* 2005;49:378–82.
13. Shih SR, Shun CT, Su DH, Hsiao YL, Chang TC. Follicular variant of papillary carcinoma: diagnostic limitations of fine needle aspiration cytology. *Acta Cytol.* 2005;49:383–6.

**Source of Support:** Nil. **Conflict of Interest:** None Declared.

**Copyright:** © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882. This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Cite this article as:** Arjun Lal Jain, Sachin Jain, Jitendra Acharya. Assessment of the Role of Fine Needle Aspiration Cytology as Diagnostic Criteria in Follicular Patterned Lesion of Thyroid. *Int J Med Res Prof.* 2017 Nov; 3(6):487-89. DOI:10.21276/ijmrp.2017.3.6.108