

# Comparative Evaluation of Intraoperative and Postoperative Parameters in Laparoscopic vs Open Appendectomy in Cases of Acute Appendicitis

Rajiv Verma<sup>1</sup>, Amit Agarwal<sup>2\*</sup>, Apoorv Goel<sup>1</sup>, Prakhar Garg<sup>2</sup>,  
Tripta S Bhagat<sup>3</sup>, Atul K Gupta<sup>3</sup>, Shalabh Gupta<sup>3</sup>

<sup>1</sup>Associate Professor, <sup>2</sup>Assistant Professor, <sup>3</sup>Professor,  
Department of General Surgery, Santosh Medical College & Hospital, Ghaziabad, Uttar Pradesh, India.

## ABSTRACT

**Background:** Appendectomy is the most common surgical procedure performed in emergency surgery. Appendectomy is still being performed by both open (OA) and laparoscopic (LA) methods as no other technique is formulated. In this study, we aimed to compare the laparoscopic procedure and the standard technique in the treatment of acute appendicitis.

**Methods:** Retrospectively collected data from 60 consecutive patients with acute appendicitis were studied. These comprised 30 patients who underwent conventional appendectomy and 30 patients treated laparoscopically. The two groups were compared for Intra operative parameters like duration of surgery, conversion, complications, and post-operative parameters like pain, requirement of pain medications, wound complications, hospital stay, any other complications, and cosmetic outcome.

**Results:** In our study Laparoscopic Appendectomy have a shorter hospital stay (Mean duration of hospital stay after surgery was 3.1 & 1.9 day in OA & LA group respectively), Operative time was significantly less in the open group (Mean duration of surgery was 71.2 minutes in OA group and 48.8 minutes in LA group). Total number of complications were fewer in the LA group with a significantly less incidence of wound infection (5 Vs 11, P <0.04).

**Conclusion:** The laparoscopic method is a safe and competent operative method in appendectomy. Postoperative

pain in our study is notably lesser in laparoscopic group as compared to open appendectomy, also post-operative complication especially wound infections are less frequent in laparoscopic group of patients. Patient's recovery from operation is also better in laparoscopic appendectomy group which includes early bowel activity, minimum hospital stay and early return to work. Overall cosmetic outcome is preferable in laparoscopic group of patients.

**Keywords:** Open Appendectomy (OA), Laparoscopic Appendectomy (LA), Hospital Cost, Appendicitis.

## \*Correspondence to:

**Dr Amit Agarwal,**  
Assistant Professor,  
Department of General Surgery,  
Santosh Medical College & Hospital,  
Ghaziabad, Uttar Pradesh, India.

## Article History:

Received: 05-12-2019, Revised: 02-01-2020, Accepted: 27-01-2020

Access this article online	
Website: <a href="http://www.ijmrp.com">www.ijmrp.com</a>	Quick Response code 
DOI: 10.21276/ijmrp.2020.6.1.048	

## INTRODUCTION

Appendicitis is the inflammation of vermiform appendix. Appendicitis is the most common cause of surgical abdomen in all age groups.<sup>1,2</sup> Persons between 10 and 19 years of age have the highest incidence of appendicitis, with males having a higher rate than females for all age groups. The lifetime risk for appendicitis has been estimated at 8.6% for males and 6.7% for females. Overall, the incidence appears to be approximately 120 per 100 000 population.

The classical features of acute appendicitis begin with poorly localized colicky abdominal pain. This is due to midgut visceral discomfort in response to appendiceal inflammation and obstruction. The pain is frequently first noticed in the periumbilical region and is similar to, but less intense than, the colic of small

bowel obstruction. Central pain abdomen is associated with loss of appetite and usually few episodes of vomiting that follow the onset of pain (Murphy). Anorexia is a useful and constant feature, particularly in paediatric age. The patient often gives a history of similar discomfort that settled spontaneously. A family history is also useful as up to one-third of children with appendicitis have a first-degree relative with a similar history.

Acute appendicitis sometimes become difficult to diagnose as it differs in the presentation in other surgical emergencies. The differential diagnosis differs according to age; in women, additional possible diagnoses are diseases of the female genital tract. In a meta-analysis review it is said that acute appendicitis is diagnosed clinically. There are many clinical methods and scoring

systems are available for diagnosis of acute appendicitis. The most broadly used scoring system is the Modified Alvarado score.<sup>6</sup> Removal of appendix (Appendectomy) is standard treatment for appendicitis. It is done preferably before development of phlegmon or after an interval of 6-8 weeks (Interval appendectomy). Since the first description of laparoscopic appendectomy by Kurt Semm<sup>2</sup> in 1983, it is being used more and more commonly. Role of laparoscopic appendectomy for the treatment of acute appendicitis is yet not clearly defined. Studies done so far have given mixed results, some favoring one or the other technique. A study at Israeli Hospital<sup>4</sup> favors open appendectomy over laparoscopic approach but another study published in American journal of surgery is in favor of laparoscopic approach.<sup>5</sup> Hence role of laparoscopic appendectomy viz a viz open appendectomy and its benefits and risks need to be clearly defined. Further studies are required to be clear about the most beneficial surgical approach to appendicitis. Present study is an effort in the same direction.

### AIMS & OBJECTIVES

The aim of this study is to evaluate comparatively laparoscopic and conventional "open" appendectomy in the treatment of acute appendicitis regarding various

1. Intra operative parameters like duration of surgery and complications.

2. Post-operative parameters like pain, requirement of pain medications, wound complications, hospital stay, and cosmetic outcome.

### MATERIALS AND METHODS

This study was done on adult patients undergoing appendectomy for appendicitis at Santosh Medical College & Hospital, Ghaziabad. This study design is a prospective one. The duration of study was taken from May 2015 to May 2016 in Department of General Surgery, Santosh Hospital & Medical College, Ghaziabad. Thirty patients of each of open appendectomy and laparoscopic appendectomy were taken in this study.

#### Statistical Analysis

All information was recorded on the standard proforma attached. Descriptive Statistical analysis was employed to describe data for frequencies, percentages, ratios, range and mean value with one standard deviation. Data were tabulated and entered in Microsoft excel.

Analysis was done with the help of IBM SPSS Statistics version 17/GenIE/Open Bug. Descriptive statistics of the variable from the data collected was carried out. Statistical analysis was performed using the chi-square test or student's t-test Fischer test as applicable. Statistical significance was defined if the p value was <0.05.

Table 1: Age distribution

Age (in years)	Open Group (N=30)		Lap Group (N=30)		p value
	Number	Percentage	Number	Percentage	
15-24	19	63.3	18	60.0	0.54* (NS)
25-35	7	23.3	10	33.3	
>35	4	13.3	2	6.7	
Mean ± SD	25.3 ± 7.8		25.3 ± 6.0		0.98**

\*Chi-square test, \*\*t-test

Table 2: Modified Alvarado Score (MAS)

MAS Score	Open Group (N=30)		Lap Group (N=30)		p value*
	Number	Percentage	Number	Percentage	
≥7	23	76.7	19	63.3	0.39 (NS)
<7	7	23.3	11	36.4	
Mean ± SD	5.2 ± 0.8		4.4 ± 0.6		0.004 (S)

\*Fisher exact test

Table 3: Duration of surgery

Duration (in minute)	Open Group (N=30)		Lap Group (N=30)		P value*
	Number	Percentage	Number	Percentage	
30 – 60	11	36.7	19	63.4	
60 – 90	12	40.0	10	33.3	0.03(S)
90 – 120	7	23.3	1	3.3	
Mean ± SD	71.2 ± 8.6		48.8 ± 8.7		0.002

\*Chi-square test

Table 4: Post-operative Visual Analogue Scale (VAS) on the day of surgery

POD 0	VAS Score			Mean ± SD	P value*
	0-3	4-6	7-10		
Open Group	7 (23.3)	17 (56.7)	6 (20.0)	7.0 ± 1.1	
Lap Group	15 (30.0)	12 (40.0)	3 (10.0)	6.0 ± 1.1	0.0001

\*Chi-square test

Table 5: Post-operative day 3

POD 3	VAS Score			Mean $\pm$ SD	P value*
	0-3	4-6	7-10		
Open Group	23 (76.7)	6 (20.0)	1(3.3)	1.9 $\pm$ 0.3	0.01(S)
Lap Group	29(96.7)	1 (3.3)	0 (0.0)	1.2 $\pm$ 0.5	

\*Chi-square test

Table 6: Post-operative Complications

Post-operative Complication	Open Group (N=30)	Lap Group (N=30)	P value
Fever	2	0	0.04
Vomiting	1	1	
Nausea	1	1	
Urinary retention	1	0	
Breathlessness	0	1	
Paralytic ileus	3	1	
Wound complication	6	2	
*Total	11	5	

\*Total number of patients with complication does not match the sum total of individual complications since some patients had more than one complication.

Table 7: Duration of hospital stay

Duration (in day)	Open Group (N=30)		Lap Group (N=30)		p value
	Number	Percentage	Number	Percentage	
1	0	0.0	10	33.3	0.001* (S)
2	2	6.7	13	43.3	
3	22	73.3	7	23.3	
4	6	20.0	0	0.0	
Mean $\pm$ SD	3.1 $\pm$ 0.5		1.9 $\pm$ 0.8		0.05

\*Chi-square test

Table 8: Post-operative cosmesis at 12 weeks

Response	Open Group (N=30)		Lap Group (N=30)		p value*
	Number	Percentage	Number	Percentage	
Very good	9	30.0	23	76.7	0.002 (S)
Good	13	43.4	4	13.4	
Satisfactory	6	20.0	2	6.6	
Poor	2	6.6	1	3.3	

\*Chi-square test

## OBSERVATIONS AND RESULTS

In this study 63.3% of OA group and 60.0% of LA group of study population belonged to age group 15 to 24 years whereas 23.3% & 33.3% of OA group and 13.3% & 6.7% of LA group of study population belonged to 25 to 35 years and 35 to 45 years respectively. But distribution of age among participants in two groups was statistically not significant ( $p>0.05$ ). Mean age of study participants was 25.3 years with 7.8 SD in OA group and 25.3 years with 6 SD in LA group but difference between mean age was statistically not significant ( $p<0.05$ ). All male and female patients were included in this study.

Table 2 show Modified Alvarado Score (MAS) in both groups. 76.7% of OA & 63.3% LA group of study participants had  $\geq 7$  MAS score. 23.3% of OA & 36.4% LA group of study participants had  $<7$  MAS score. Difference between score of study participants of two groups was statistically not significant ( $p>0.05$ ). Mean MAS score was 5.2 with 0.8 SD in OA group and 4.4 with 0.6 SD in LA group and difference was statistically significant ( $p<0.05$ ).

Pre-operative USG was suggestive of acute appendicitis is 60% (18/30) of OA group & 73.3(22/30) of LA group of patients. In 9 & 8 patients respectively in OA and LA group pre-operative USG was not done due to non-availability of USG at that time. In 3 patients in PA group pre-operative USG was not suggestive of acute appendicitis.

Table 3 shows that surgery was completed in 30 to 60 minutes in 36.7% of OA group & 63.4% of LA group of study participants. In 60 to 90 minutes 40.0% of OA group & 33.3% of LA group of study participants completed surgery. In 23.3% cases of OA & 3.3% cases of LA group have surgery time 90 to 120 minutes. This difference in time of surgery completion between OA & LA group was statistically significant ( $p<0.05$ ). Mean duration of surgery was 71.2 minutes with 8.6 SD in OA group and 48.8 minutes with 8.7 SD in LA group and difference was statistically significant ( $p<0.05$ ).

Table 4 shows comparison of post-operative VAS score between OA and LA group on the day of surgery. 23.3% of OA & 30.0% of

LA group of cases had VAS score 0 to 3. 56.7% of OA & 40.0% of LA group of cases have VAS score 4 to 6 and 20.0% of OA & 10.0% of LA group cases had VAS score 7 to 10. This difference in VAS score between these groups was statistically not significant ( $p>0.05$ ). Mean VAS score on the day of surgery of OA group was  $7.0 \pm 1.1$  & of LA group was  $6.0 \pm 1.1$  and difference was statistically significant ( $p<0.05$ ).

Table 5 shows comparison of post-operative VAS score between OA and LA group after day 3 of surgery. 76.7% of OA & 96.7% of LA group of cases had VAS score 0 to 20.0% of OA & 3.3% of LA group of cases had VAS score 4 to 6 and 3.3% of OA cases had VAS score 7 to 10. This association of VAS score between these groups was statistically not significant ( $p>0.05$ ). Mean VAS score on POD3 of OA group was  $1.9 \pm 0.3$  & of LA group was  $1.2 \pm 0.5$  and difference was statistically significant ( $p<0.05$ ). All patients were given diclofenac sodium 75mg by intramuscular route twice a day & switched to oral later. Extra analgesic was required in 23.3% of OA & 10.0% of LA group of study participants during surgery. This difference is statistically not significant ( $p>0.05$ ).

In OA group 2 patients had fever post operatively. Both patients recovered without any intervention. Equal number of patients in both groups suffered from nausea and vomiting. Antiemetic were given to the patient to subside the symptoms. 1 patient was catheterized postoperatively for urinary retention in OA group. 6 patients in OA group suffered from paralytic ileus, out of this, four patients recovered within 2 days and remaining patients within 4 days. In all individuals only conservative approach was sufficient. In LA, 1 patient suffered paralytic ileus, same treatment was given. Total of 6 patients in OA group had postoperative wound complication as compared to only 2 patients in LA group. 3 patients did not need any additional measures as due to superficial wound infection. 2 patients had leakage of seroma from wound site, which was relived with removal of suture. Remaining one patient had slough on operative site which was debrided and was given antibiotic. Both patient in LA group had port site infection which resolved with dressing.

Table 7 shows that 73.3% of OA & 23.3% LA group of participants stayed for 3 days after surgery and 6.7% of OA & 43.3% of LA group of participants stayed in hospital for 2 days after surgery. 20.0% study participants of OA group stayed for 4 days after surgery. This difference between duration of post-operative stay at hospital was statistically significant ( $p<0.05$ ). Mean duration of hospital stay after surgery was 3.1 & 1.9 day in OA & LA group respectively and difference was statistically significant ( $p<0.05$ ).

Table 8 shows that 30.0% of OA & 83.3% of LA group of study participants had an excellent cosmetic end result. Patients were asked to grade the cosmetic outcome as very good, good, satisfactory or poor at 12 weeks follow up. 23.3% of OA & 3.3% of LA group of participants have satisfactory cosmetic end result. This difference was statistically significant ( $p<0.05$ ).

## DISCUSSION

Appendiceal disease is a frequent reason for emergency hospital admission, and appendectomy is one of the most common emergency procedures performed in contemporary medicine. Although in this advanced world, we have much radiographic imaging system available with modern diagnostic tools, laboratories and accurate investigations, still the diagnosis of acute appendicitis is made by clinical parameters.

In this study we considered total 60 patients out of which 30 in OA & 30 in LA group.

In both groups, the participants belonged to 15 to 25 years age group which constitutes about 60% of total and shows that the younger age groups are frequent participants. This data establish that appendicitis is the disease of young age group which is a worldwide fact. Mean age of participants was  $25.3 \pm 7.8$  years in OA and  $25 \pm 6.0$  years in LA & the difference between mean age of both the groups was statistically not significant as shown in table 1. It implies that distribution of patients regarding age in two groups was equitable. Similar statistically non-significant results of mean age were found by Thakre S et al.<sup>7</sup>, and Khan MN et al.<sup>8</sup>

There were 14 female & 16 male in OA group and 16 female & 14 male in LA group. Male to female ratio was 1.14:1 in open appendectomy and 0.87:1 in Laparoscopic appendectomy group which shows that female has higher number in LA group in comparison to OA group.

The Modified Alvarado Scoring System (MASS) has been reported to be a cheap and quick diagnostic tool in patients with acute appendicitis. Modified Alvarado Score (MAS)  $\geq 7$  was observed in present study in 76.7% of OA & 63.3% LA group of patients (Table 2). In 23.3% of OA & 36.7% LA group of study participants MAS score was  $< 7$ . Results of present study are comparable with the study done by Verma M et al<sup>9</sup>, and Sonawane RS et al.<sup>10</sup>

USG is being increasingly used in diagnosing acute appendicitis. In our study only 60% of patients in OA group & 73.3% in LA group, had pre-operative USG findings suggestive of acute appendicitis.

In OA group, the operative duration was approximately 90 minutes as compared to LA group, in which majority of patients were operated in less than 60 minutes. Mean duration of surgery was 71.2 minutes in OA group and 48.8 minutes in LA group as shown in Table 3. Open appendectomy took much higher time than the laparoscopic appendectomy and difference was statistically significant. This result is similar with the results of study done by Mwero BJ et al.<sup>11</sup> and Puser JG et al.<sup>12</sup> where duration of surgery was significantly higher in OA group than LA group.

We observed visual analogue scale (VAS) for postoperative pain on the day of surgery, and 3<sup>rd</sup> post-operative day after surgery where VAS score is between 1 to 10. Every patient was assessed first for VAS scoring before giving analgesics.

On the day of operation mean VAS score of OA group is  $7.0 \pm 1.1$  and in LA group is  $6.0 \pm 1.1$ . This difference is statistically significant as shown in table 4. Lesser post-operative pain after laparoscopic appendectomy as compared to open appendectomy has been documented by Chung RS et al<sup>13</sup>, Garbutt JM et al<sup>14</sup>, Golub R et al<sup>15</sup> and Thakre S et al.<sup>7</sup>

Extra analgesia was used on the basis of VAS scoring, clinical assessment & patient demand. Patients were given extra analgesia if VAS  $> 5$ .

In present study more patients experienced post-operative complication in OA group compared to LA group (Table 6). One patient in OA group experienced retention of urine which could be due to the effect of spinal anesthesia. One patient in LA group complained breathlessness in post-operative period which subsided in few hours, the cause of it could not be ascertained. Patients had complications as fever, nausea, vomiting, slightly more in OA group compared to LA group. Study done by

Chitumalla et al<sup>16</sup> had less wound infection following laparoscopic appendectomy as compared to open appendectomy. The difference between two groups was statistically significant regarding post-operative wound infection. Postoperative complications were more in patients underwent open appendectomy as compared to laparoscopic appendectomy in a study done by Guller et al.<sup>17</sup>

In our study Laparoscopic appendectomy group patients had earlier return of bowel activity as compared to those in open appendectomy group. 80% patients in LA group had post-operative bowel movements on POD1 as compared to only 50% in OA group. As shown by Ortega AE et al.<sup>18</sup> and Hellberg A et al<sup>19</sup> show similar result of return of early bowel activity in LA group as compared to OA group.

As shown in table 7, large majority of patients (23/30) in LA group were discharged by 2<sup>nd</sup> post-operative day. In contrast in OA group maximum patients (22/30) were discharged on 3<sup>rd</sup> post-operative day. Mean duration of hospital stay after surgery was 3.1 & 1.9 day in OA & LA group respectively. Significantly longer post-operative hospital stays in LA group as compared to OA group in our study is in agreement with the studies done by Golub R et al.<sup>15</sup> and Moberg AC et al.<sup>20</sup>. Primarily less postoperative pain accounts for reduced hospital stay, minimum requirement of extra analgesia, early bowel sounds, minimum postoperative complications and early starting of diet. For earlier return to normal activity minimum hospital stay is essential. This decreases hospital cost & burden on health care system. But studies done by Chung RS et al<sup>13</sup>, Garbutt JM et al<sup>21</sup>, Ignacio RC et al<sup>22</sup> found no statistically significant difference in hospital stay after surgery between two groups.

Appendicitis is disease of predominantly young people. Since significant number of patients undergoing appendectomy are young females, cosmesis is an important factor. In our study we asked the patients at 12 weeks follow up to grade the cosmetic outcome of operative scar as excellent, good, satisfactory & poor. Patients in LA group graded their cosmetic outcome significantly better in LA group than in OA group (p value .002). 83.3% patients reported their cosmetic result as 'excellent' in LA group as compared to only 30% in OA group (Table 8). Study done by Chitumalla et al<sup>16</sup> showed significant difference in cosmesis among two groups. Patients underwent Laparoscopic appendectomy in their study group had excellent cosmesis.

## CONCLUSION

Laparoscopic appendectomy is a safe and promising procedure which needs an upper degree of technical skill on the part of a surgeon. Laparoscopic appendectomy is a hopeful alternate way to conventional appendectomy.

Appendicitis is a clinical diagnosis. Post-operative pain in our study is significantly fewer in laparoscopic groups as compared to open appendectomy. The post-operative complication most importantly wound infections are less frequent in laparoscopic group of patients.

Recovery of patients from operation is also better in laparoscopic group including early bowel motion, shorter hospital stay and early return to work. Overall cosmetic outcome is preferable in laparoscopic group of patients. Thus, laparoscopic appendectomy is an accepted and reliable procedure. Availability of laparoscopic set up and expertise of surgeon is a

challenge. Though conventional appendectomy is widely used procedure for appendectomy, laparoscopic appendectomy is proving to be better procedure of choice in cases of acute appendicitis.

## REFERENCES

1. Addiss DG, Shaffer N, Fowler BS, Tauxe R. The epidemiology of appendicitis and appendectomy in the United States. *Am J Epidemiology*. 1990;132:910–25.
2. Semm K. Endoscopic appendectomy. *Endoscopy*. 1983; 15: 59–64.
3. Khalil J, Muqim R, Rafique M, Khan M. Laparoscopic versus open appendectomy: A comparison of primary outcome measures. *Saudi J Gastroenterol* 2011; 17:236-40.
4. The American Journal of Surgery, Volume 169, Issue 2, Pages 208–13.
5. Herrington JL Jr. The vermiform appendix: its surgical history. *Contemp Surg*. 1991; 39:36-43.
6. Kalan M, Talbot D, Cunliffe WJ, Rich AJ. Evaluation of the modified Alvarado score in the diagnosis of acute appendicitis: a prospective study. *Ann R Coll Surg*. 1994;11:418–9.
7. Thakre S, Singh VM, Mudgal MM, Kushwah N, Gupta A. A Comparative Study between Laparoscopic Appendectomy and Conventional Open Appendectomy. *Sch. J. App. Med. Sci.*, 2014; 2(5F):1909-12.
8. Khan MN, Fayyad T, Cecil TD, Moran BJ. Laparoscopic versus open appendectomy: the risk of postoperative infectious complications. *JLS*. 2007;11:363–7.
9. Verma M, Senior MS, Surgery G, Pt BD, Surgery G, Post-PBDS, et al. Comparison of Alvarado And Ripasa Scoring Systems in Diagnosis of Acute Appendicitis. 2015;55–7.
10. Sonawane RS, Jatkar GL, Chaudhari MS. Correlation of Alvarado score for acute appendicitis with pathological acute appendicitis. 2016;3(3):1451–5.
11. Mweru BJ. Laparoscopic Appendectomy Versus Open Appendectomy At Kenyatta National Hospital a Dissertation Submitted in Part Fulfilment for the Degree of Master of Medicine (Surgery), University of Nairobi. 2003.
12. Puser Jochanan G; Greenberg Dan. Laparoscopic vs open appendectomy: Results of a retrospective comparison in Israel Hospital. *Israel Medical Association Journal*; IMAS (Siraal; Feb 2002; 4; (2): p 91 – 4.
13. Chung RS, Rowland DY, Li P, et al. A meta-analysis of randomized controlled trials of laparoscopic versus conventional appendectomy. *Am J Surg*. 1999;177:250–6.
14. Garbutt JM, Soper NJ, Shannon WD, et al. Meta-analysis of randomized controlled trials comparing laparoscopic and open appendectomy. *Surg Laparosc Endosc*. 1999;9:17–26.
15. Golub R, Siddiqui F, Pohl D. Laparoscopic versus open appendectomy: a meta-analysis. *J Am Coll Surg*. 1998; 186: 545–53.
16. Chitumalla P, Reddy AST, Reddy SV. Comparative study of open versus laparoscopic appendectomy outcome in our local area. *J. Evid. Based Med. Healthc*. 2016;3(39):1931-37.
17. Guller U, Hervey S, Purves H, et al. Laparoscopic versus open appendectomy: outcomes comparison based on a large administrative database. *Ann Surg*. 2004;239:43–52.
18. Ortega AE, Hunter JG, Peters JH, Swannstrom LL, Schirmer B. Laparoscopic Appendectomy Study Group. A prospective,

randomized comparison of laparoscopic appendectomy with open appendectomy. *Am J Surg.* 1995; 169:208–13.

19. Hellberg A, Rudberg C, Kullman E, Enochsson L, Fenyö G, Graffner H, Hallerbäck B, Johansson B, Anderberg B, Wenner J, Ringqvist I, Sörensen S. Prospective randomized multicentre study of laparoscopic versus open appendectomy. *Br J Surg.* 1999;86:48–53.

20. Moberg AC, Montgomery A. Appendicitis: laparoscopic versus conventional operation: a study and review of the literature. *Surg Laparosc Endosc.* 1997;7:459–63.

21. Garbutt JM, Soper NJ, Shannon WD, Botero A, Littenberg B.. Meta-analysis of randomized controlled trials comparing laparoscopic and open appendectomy. *Surg Laparosc Endosc.* 1999;9:17–26.

22. Dr Tom Lomax. The pros and cons of inpatient and outpatient care < Japan Today Japan News and Discussion Sep. 15, 2015.

**Source of Support:** Nil. **Conflict of Interest:** None Declared.

**Copyright:** © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882. This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Cite this article as:** Rajiv Verma, Amit Agarwal, Apoorv Goel, Prakhargarg, Tripta S Bhagat, Atul K Gupta, Shalabh Gupta. Comparative Evaluation of Intraoperative and Postoperative Parameters in Laparoscopic vs Open Appendectomy in Cases of Acute Appendicitis. *Int J Med Res Prof.* 2020 Jan; 6(1): 200-05. DOI:10.21276/ijmrp.2020.6.1.048