

Study of Short Stature in Children with Special Reference to Etiology and Management: An Hospital Based Study

Khurshida Khan¹, Geetanjali Shrivastava^{2*}, N. Saravanan², Uma Raju³

¹Assistant Professor, Department of Pediatric Medicine, Government PDU Medical College, Churu, Rajasthan, India.
(Ex fellow, Department of Neonatology, Nice Hospital for Women, Newborn & Children, Hyderabad)

²Consultant Neonatology, Department of Neonatology, Nice Hospital for Women, Newborn & Children, Hyderabad, India.

³Head, Clinical and Academic Services, Nice Hospital for Women, Newborn & Children, Hyderabad, India.

ABSTRACT

Background: Short stature is a common problem in children globally, especially in developing countries. Idiopathic Short Stature (ISS) is considered when no causative disorder can be identified. The aim of this study is to find out the etiological profile of short stature, its management and the effect of intervention on stature.

Materials & Methods: This is a hospital based prospective study done on all indoor patients with age 2 months to 2 years. Patients with inadequate follow up and Neonates and infants < 2 months were excluded in our study. All children with age < 12 years if they presented with height < 3rd percentile for age.

Results: Out of 2600, 260 cases of short stature. So the incidence of short stature in the present study was 10%. Maximum incidence of short stature is found to be among the age group of 1-5 years which is 51.9%. Male: Female ratio was 0.95. Bone age was normal (equal to chronological age) in 32.6% of patients, while it was abnormal (<chronological age) in 67.4% patients. Incidence of proportionate short stature in this study is 88% and disproportionate is 12%. In etiological profile of short stature Pathological (65%) is more common than Physiological (35%). In Physiological short stature

category, among admitted patients- Constitutional (63.4%) is more common than familial (36.6%).

Conclusion: We concluded that incidence of short stature was 10%, mostly occurred in 1-5 years of age and etiological profile of short stature Pathological (65%) is more common than Physiological (35%).

Keywords: Short Stature, Etiology, Physiological, Pathological, Growth.

*Correspondence to:

Dr. Geetanjali Shrivastava,
Consultant Neonatology,
Department of Neonatology, Nice Hospital for Women,
Newborn & Children, Hyderabad, India.

Article History:

Received: 26-09-2020, Revised: 28-10-2020, Accepted: 23-11-2020

Access this article online

Website: www.ijmrp.com	Quick Response code 
DOI: 10.21276/ijmrp.2021.7.1.012	

INTRODUCTION

Short stature is a common problem in children globally, especially in developing countries. When compared with well-nourished and genetically relevant population, short stature is defined as height or length below 3rd percentile for that age and gender. Statistically, this refers to children who are shorter than 97% of their age and gender matched peers.¹

Although a small body size can simply be within the normal ranges, it may also indicate various possible medical conditions. Causes of short stature are diverse, it may be genetic, metabolic, chronic systemic illness or simple environmental issues such as reduced food availability.¹

Almost any chronic disease can cause short stature such as renal disease, malignancy, pulmonary disease, Cystic Fibrosis (CF), cardiac disease etc. Coeliac disease is a prime example of a remediable cause of short stature, especially in younger children.²

Nutritional deprivation and therapies like glucocorticoids, chemotherapeutic drugs, radiotherapy can result in short stature. Common endocrinological causes of short stature include hypothyroidism, hypopituitarism (isolated GHD or multiple anterior pituitary hormone deficiencies), hypercortisolism and classical Laron syndrome. All these are characterized by being overweight-for-height.² Short stature may also be seen with severe Intrauterine Growth Retardation (IUGR) or children born Small for Gestational Age (SGA) and in large number of dysmorphic syndromes. Idiopathic Short Stature (ISS) is considered when no causative disorder can be identified. Anybody change or abnormality needs prompt professional medical advice for medical investigation and diagnosis.³ The aim of this study is to find out the etiological profile of short stature, its management and the effect of intervention on stature.

MATERIALS & METHODS

This is a hospital based prospective study done on all indoor patients with age 2 months to 2 years. Patients with inadequate follow up and Neonates and infants < 2 months were excluded in our study.

Selection Criteria for Short Stature in Children

All children with age < 12 years if they presented with height < 3rd percentile for age on:

National Centre for Health Statistics-Centre for Disease Control (2 yr. to 12 yr.) growth curve

World Health Organisation (2 months to 2 yr.) curves are chosen as reference.

Anthropometry

Height measurement is performed with child standing erect

without shoes with heels together, arms resting at the side of the body, heels buttocks occiput touching the wall and head positioned in the Frankfurt plane (>2 Years) by Stadiometer.

Length measurement is performed with the child lying supine (2 months – 2 years) by Infantometer⁴. Upper segment (sitting height) / lower segment is taken for all patients.

Weight performed on electronic machine.

Nutritional evaluation of patients < 5 years, weight / height < -2SD were considered as malnourished according to World Health Organisation criteria.

For patients >5 years, BMI is calculated and those with BMI <5 pc was considered as malnourished according to Center for Disease Control criteria & graph.

Table 1: Age & Sex- Wise Distribution

Age	Male	Female	No.	Percentage
2M –1Y	7	8	15	5.7%
1Y – 5Y	65	70	135	51.9%
5Y-9Y	38	32	70	26.9%
9Y-12Y	17	23	40	15.5%
Total	127	133	260	-
Percentage	48.8%	51.2%	-	100%

Table 2: Etiological Profile:

Etiology		Number	Percentage
Physiological (N=90)	Constitutional short stature	52	63.4%
	Familial	38	36.6%
Pathological (N=170)	Respiratory system	27	10.3%
	Cardiovascular system	21	8.1%
	Renal system	17	6.6%
	Hematological system	71	27.3%
	Genetic syndrome	15	5.7%
	Endocrine system	15	5.7%
	Gastrointestinal system	6	2.4%
	Orthopedic disorder	8	3%
	Metabolic disorders	19	7.4%
	Central nervous system	14	5.3%

Table 3: Short Stature Type:

Proportionate	230(88%)
Disproportionate	30(12%)


RESULTS

Total admissions during study period were 2600 cases. Out of 2600, 260 cases of short stature. So, the incidence of short stature in the present study was 10%. Maximum incidence of short stature is found to be among the age group of 1-5 years which is 51.9%. Male: Female ratio was 0.95 (table 1).

In etiological profile of short stature Pathological (65%) is more common than Physiological (35%). In Physiological short stature category, among admitted patients- Constitutional (63.4%) is more common than familial (36.6%). In Pathological short stature category, incidences among admitted patients are as follows- Malnutrition (16.5%), respiratory system (10.3%), cardiovascular

system (8.1%), renal system (6.6%), endocrinological (5.7%), haematological (27.3%), Metabolic disorders (7.4%) and Genetic syndromes (5.7%) (table 2).

Bone age was normal (equal to chronological age) in 32.6% of patients, while it was abnormal (<chronological age) in 67.4% patients (figure 1). Incidence of proportionate short stature in this study is 88% and disproportionated is 12% (table 3). Patients of physiological short stature, have low growth velocity but in their target range. Patients of pathological short stature category, Early recognition of growth abnormality and early intervention maintains adequate growth velocity.



Graph 1: Bone age analysis

DISCUSSION

In the present study the incidence of short stature among total hospital admissions is 10%. A study done by Colaco et al⁵, incidence of short stature was 5.6% and Khadgawat et al⁶ found incidence of short stature among total hospital admissions was 7%. In the present study, the incidence of short stature in male:female is 0.95. Chowdhury SP et al⁷ found was 0.80 and Sultan M et al⁸ observed 1.03 male to female ratio. The incidence of short stature was maximum in 1-5 yrs which is 54%, which was compatible with Sultan M et al⁸ found 43.5%. In etiological profile of short stature Pathological (65%) is more common than Physiological (35%). Variou study done by Sultan M et al⁸, Garg P⁹ and Chowdhury SP et al⁴⁵ found the percentage of pathological short stature (62% 75% & 82% respectively) is greater than physiological (38%, 25% & 18% respectively). As per this study, the incidence of constitutional short stature (22%) is greater than familial (12.7%). Moayeri et al¹⁰ showed that the incidence of constitutional short stature (22%) is greater than familial (14%).

The incidence of short stature in the present study with patients of malnutrition is 16.5% and in the study done by Sultan M et al⁸ is 9.8%. The difference can be attributed to higher rate of admission of patients of low socio-economic level with greater occurrence of malnutrition among them in our tertiary care centre. The incidence of chronic anemia in the present study is 17.3% as this tertiary care centre holds thalassemia clinics which explains the difference with the study by Sultan M et al⁸ where incidence is 1.9%.

In the present study, the incidence of proportionate short stature is 88 & disproportionate is 12%. Chowdhury SP et al⁴⁵ found the incidence of proportionate short stature is 48 & disproportionate is 52%. In this study, inadequate growth velocity was found among asthma patients with poor control which is similar to study done by Mc Cowan 1998¹ which showed decreased growth velocity in patients with high dose steroids and requiring frequent hospital visits. Among pts. of acyanotic CHDs control of heart failure & proper nutrition have adequate GV. While pts. of cyanotic CHDs have invariably poor GV except after TGA surgical correction. In the study done by Soliman AT¹¹ size of L-R shunt, hemodynamics, hypermetabolic state are all factors determining growth velocity with normal growth velocity by control of heart failure.

CONCLUSION

We concluded that incidence of short stature was 10%, mostly occurred in 1-5 years of age and etiological profile of short stature Pathological (65%) is more common than Physiological (35%).

REFERENCES

1. KE Elizabeth, Nutrition and child development, 4th ed, sec 2:64.
2. Kleigman & Behrman, Nelson Textbook of Paediatrics, 18th edition, volume 1, ch-12, pg-60.
3. O.P. Ghai, Essential Pediatrics, 7th edition, pg-03
4. ML Kulkarni. Clinical method in Pediatrics, volume 2 part A,pg-89.
5. Colaco P, Desai M, Choksi CS. Short stature in Indian children: the extent of the problem. Indian J Pediatr. 1991;58 Suppl 1:57-8.
6. Khadgawat R, Dabadghao P, Mehrotra RN, Bhatia V. Growth charts suit- able for evaluation of Indian children. Indian Pediatr. 1998;35:859-65.
7. Chowdhury SP, Sarkar TK, Haldar D, Taraphdar P, Naskar TK, Sarkar GN. Short stature in children: Experience from a tertiary care hospital in Kolkata, India. The Health. 2011;2(4):139-42.
8. Sultan M, Afzal M, Qureshi SM, Aziz S et al. Etiology of short stature in children. J Coll Physicians Surg Pakistan. 2008;18(8):493-7.
9. Garg P. Short stature in Indian children: experience from a community level hospital. Sri Lanka J Child Health. 2004;34:84-8.
10. Moayeri H, Aghighi Y. A prospective study of etiology of short stature in 426 short children and adolescents. Arch Iranian Med 2004; 7:23-7.
11. Soliman AT, Madkour A, Galil MA, El Zalabany M, Aziz SM, Ansari BM. Growth parameters and endocrine function in relation to echocardiographic parameters in children with ventricular septal defect without heart failure. J Trop Pediatr 2001 Jun;47(3):146-52.

Source of Support: Nil.

Conflict of Interest: None Declared.

Copyright: © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882. This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article as: Khurshida Khan, Geetanjali Shrivastava, N. Saravanan, Uma Raju. Study of Short Stature in Children with Special Reference to Etiology and Management: An Hospital Based Study. Int J Med Res Prof. 2021 Jan; 7(1): 42-44. DOI:10.21276/ijmrp.2021.7.1.012