

Assessment of Osteoporosis Knowledge among Female Medical Interns And Students at Tabuk University

Wejdan Abdulrahman Alqarni^{1*}, Ashwaq Ali Absalem¹, Nadiyah Abdullah Almountashiri¹,
Afrah Dhaifallah Alanazi², Wafa Hammad Albalawi², Elaf Ibrahim Alanzi²,
Norah Abdulrahman Alfuraydi², Afaf Atallah Albalawi², Abdulaziz Fahad Alsuwayh³,
Fahad Abdullah Alqahtani³

¹Medical Intern, Ministry of Health, Kingdom of Saudi Arabia.

²Medical Student, Ministry of Health, Kingdom of Saudi Arabia

³Internal Medicine Resident, Tabuk, Saudi Arabia.

ABSTRACT

Introduction: Osteoporosis is a common disease that is characterized by low bone mass and skeletal fragility, leading to higher risk of fracture. The disease is medically diagnosed as having reduced bone mineral density that is 2.5 standard deviations below the adult peak mean. Approximately 34 % of Saudi female are osteoporotic, to treat and prevent osteoporosis our health care providers should have adequate level of knowledge about osteoporosis.

Methods: This study was carried out among medical interns at Universities of Tabuk, Saudi Arabia. A pre-tested self-administered questionnaire was used to assess knowledge, beliefs, and practices towards osteoporosis. Osteoporosis Knowledge Assessment Tool (OKAT), a validated tool was used. This consists of 20 questions to assess knowledge.

Results: Rate was 100 % with a total 81 female students from the University of Tabuk. A 45.6 % of the participants were medical interns, 32% were fifth year medical students and 22.4% were sixth year medical students. 83.9% of the participants correctly identify the relation between the age and osteoporosis, 67.9% knew that smoking is one of osteoporosis risk factors, and 46.9% correctly consider present of family

history of osteoporosis as a risk factor. 55.9% participants felt eating calcium rich food was difficult. 87.7% felt that if they had osteoporosis that it would change their whole life.

Conclusion: Majority of participants had a acceptable level of knowledge on osteoporosis, Steps such as education about osteoporosis risk factors and preventing methods are required to improve the health states of the community.

Keywords: Osteoporosis, Postmenopausal, Tabuk, Saudi Arabia.

*Correspondence to:

Wejdan Abdulrahman Alqarni,
Medical Intern, Ministry of Health, Kingdom of Saudi Arabia.

Article History:

Received: 08-11-2017, Revised: 04-12-2017, Accepted: 21-12-2017

Access this article online

Website: www.ijmrp.com	Quick Response code 
DOI: 10.21276/ijmrp.2018.4.1.046	

INTRODUCTION

Osteoporosis, a chronic, progressive disease of multifactorial etiology. It has been most frequently recognized in elderly white postmenopausal women. Osteoporosis is a common disease that is characterized by low bone mass and skeletal fragility, leading to higher risk of fracture, specially at the spine, hip, wrist, humerus and pelvis.¹ The World Health Organization has defined diagnostic thresholds for low bone mass and osteoporosis based upon bone mineral density (BMD) measurements compared with a young adult reference population (T-score). The disease is medically diagnosed as having reduced bone mineral density that is 2.5 standard deviations below the adult peak mean.² They may be diagnosed as an incidental finding on chest or abdominal radiographs. The clinical manifestations of symptomatic vertebral fractures include pain and height loss. Men may also have symptoms of disorders that are known to cause

osteoporosis, such as hypogonadism, malabsorption, and hyperparathyroidism.

An estimated prevalence of osteoporosis showed that about 34% of healthy Saudi women, and 30.7% of men, aged between 50-79 years are osteoporotic, which expected to increase further.³

For the prevention and treatment of osteoporosis, it is vital that health care providers have adequate knowledge about osteoporosis to ensure that they can effectively treat individuals with this disease.

The awareness of medical students about osteoporosis were studied in many studies, a lot of studies shown that medical students know the definition of osteoporosis, but lack knowledge of its complications and preventive measures.⁴

In this study we are aiming to assess the level of medical student's awareness about osteoporosis.

METHODOLOGY

This study was carried out among medical interns at Universities of Tabuk, Saudi Arabia. A pre-tested self-administered questionnaire was used to assess knowledge, beliefs, and practices towards osteoporosis. Osteoporosis Knowledge Assessment Tool (OKAT), a validated tool was used 5. This consists of 20 questions to assess knowledge about osteoporosis.

Ethical Considerations



Study was explained to participants and informed consent was taken from the participant.

Statistical Analysis

The collected Data were entered and analyzed using the Statistical Package for Social Sciences (SPSS) statistical program version 19.

Table1: Descriptive characteristics of the participants

		Number	Percentage
Gender	Female	81	100%
Smoking	Non Smoker	1	1.2%
	Smoker	1	1.2%
	Ex-Smoker	79	97.5%







Table 2: Assessment knowledge about osteoporosis symptoms.

Osteoporosis usually causes symptoms (e.g. pain) before fractures occur.			
	Agree	Disagree	Don not know
Interns	48.6%	45.9%	5.4%
6th year	55.5%	33.3%	11.1%
5th year	61.5%	23%	15.3%

Table 3: Assessment knowledge about osteoporosis fracture risk.

Osteoporosis leads to an increased risk of bone fractures.			
	Agree	Disagree	Don not know
Interns	18.9%	45.9%	35.1%
6th year	55.5%	10.8%	10.8%
5th year	26.9%	42.3%	30.7%

RESULTS

The response rate was 100 % with a total 81 female students from the University of Tabuk. A 45.6 % of the participants were medical interns, 32% were fifth year medical students and 22.4% were sixth year medical students. 97.5% were non-smoker, 1.2% smoker and 1.2% ex-smoker. (Table 1)

As shown in figure 1, 83.9% of the participants correctly identify the relation between the age and osteoporosis, 67.9% knew that smoking is one of osteoporosis risk factors, and 46.9% correctly consider present of family history of osteoporosis as a risk factor.

In figure 2, age and family history of osteoporosis were mostly identified by fifth year medical students, where smoking mostly identified by sixth year medical students.

Perceptions, towards benefits of calcium intake revealed that 55.9% participants felt eating calcium rich food was difficult, and 9.8% felt that calcium rich foods are too expensive. (Figure 3)

On assessing the perceived seriousness of osteoporosis, 87.7% felt that if they had osteoporosis that it would change their whole life and would make daily activities more difficult.

As shown in Table 2, approximately 50% and more of female medical students and interns belief that osteoporosis usually causes symptoms. Only 18.9% and 26.9% of medical interns and 5th year medical students aware that fracture is one of osteoporosis complication. (Table 3)

DISCUSSION

The aim of the study was to assess knowledge, beliefs and practices regarding osteoporosis among female medical interns and students at Tabuk University. Knowledge about osteoporosis preventive and its risk factors are important in preventing and minimizing its complication. However, most of study participants truly identify most of osteoporosis risk factors, including and not limited to advanced age, smoking and present of family history of osteoporosis. Low calcium intake is associated with increased osteoporosis risk. Optimal calcium intake is important to reach optimal peak bone mass, which is protective against osteoporosis and its complication, 44.1% of our study participants believe that eating calcium rich food is difficult.⁶

CONCLUSIONS

Majority of participants had an acceptable level of knowledge on osteoporosis. Steps such as education about osteoporosis risk factors and preventing methods are required to improve the health states of the community.

REFERENCES

1. Lips P, van Schoor NM. Quality of life in patients with osteoporosis. *Osteoporos Int* 2005;16:447e55.
2. Kanis JA, Melton LJ, Christiansen C, Johnston CC, Khaltav N. Perspective: the diagnosis of osteoporosis. *J Bone Mineral Res* 1994;9:1137e41.
3. Sadat-Ali M, Al-Habdan IM, Al-Turki HA, Azam MQ. An epidemiological analysis of the incidence of osteoporosis and osteoporosis-related fractures among the Saudi Arabian population. *Ann Saudi Med*. 2012 Nov-Dec; 32(6):637-41.
4. S. Eyigor, H. Karapolat, B. Durmaz Medical students' knowledge of osteoporosis in Ege University Faculty of Medicine. *Med Teach*, 13 (2009 Jan), p. 1
5. Winzenberg TM, Oldenburg B, Freuden S, Jones G. The design of a valid and reliable questionnaire to measure osteoporosis knowledge in women: the Osteoporosis Knowledge Assessment Tool (OKAT) *BMC Musculoskelet Disord*. 2003;4:17. doi: 10.1186/1471-2474-4-17.
6. Hu JF, Zhao XH, Jia JB, Parpia B, Campbell TC. Dietary calcium and bone density among middle-aged and elderly women in China. *Am J Clin Nutr*. 1993 Aug; 58(2):219-27.

Source of Support: Nil.

Conflict of Interest: None Declared.

Copyright: © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882.

This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article as: Wejdan Abdulrahman Alqarni, Ashwaq Ali Absalem, Nadiyah Abdullah Almountashiri, Afrah Dhaifallah Alanazi, Wafa Hammad Albalawi, Elaf Ibrahim Alanzi, Norah Abdulrahman Alfuraydi, Afaf Atallah Albalawi, Abdulaziz Fahad Alsawayh, Fahad Abdullah Alqahtani. Assessment of Osteoporosis Knowledge among Female Medical Interns And Students at Tabuk University. *Int J Med Res Prof*. 2018 Jan; 4(1):236-39. DOI:10.21276/ijmrp.2018.4.1.046