

A Study of Prevalence of Oral Lichen Planus in a Known Population: A Cross-Sectional Hospital Based Study

Prasoon Soni¹, Ekta Soni^{2*}, Jitendra Acharya³

¹Assistant Professor, Department of Skin & V.D., S. P. Medical College, Bikaner, Rajasthan, India.

^{2*}Senior Demonstrator, Department of Physiology, S. P. Medical College, Bikaner, Rajasthan, India.

³Senior Demonstrator, Department of Dentistry, S.P. Medical College, Bikaner Rajasthan, India.

ABSTRACT

Background: Oral lichen planus (OLP), the mucosal equivalent of cutaneous lichen planus, presents frequently in the fourth decade of life. Vesicle and bulla are seen in the oral lesions of lichen planus. Hence; we conducted the present study to assess the prevalence of OLP in a known population of Samba region, Jammu.

Materials & Methods: The present study included evaluation of prevalence rate of OLP. A total of 200 subjects were included in the present study through a joint study of department of skin and V.D. and department of dentistry, S.P. Medical College Bikaner. Detailed demographic details and clinical details of all the patients were recorded. Mouth mirror and probe were used for making the clinical examination of the patients. All the results were recorded on excel sheet.

Results: Out of total 200 subjects, 120 were males and remaining were females. OLP was found to be present in 2 males and 1 female. Prevalence rate of OLP was found to be 1.5 percent.

Conclusion: OLP occurs in a significant proportion of population.

Keywords: Oral Lichen Planus, Prevalence.

*Correspondence to:

Dr. Ekta Soni,
Senior Demonstrator,
Department of Physiology,
S. P. Medical College, Bikaner, Rajasthan, India.

Article History:

Received: 30-09-2018, Revised: 22-10-2018, Accepted: 17-11-2018

Access this article online	
Website: www.ijmrp.com	Quick Response code 
DOI: 10.21276/ijmrp.2018.4.6.059	

INTRODUCTION

Oral lichen planus (OLP), the mucosal counterpart of cutaneous lichen planus, presents frequently in the fourth decade of life and affects women more than men in a ratio of 1.4:1. The disease affects 1-2% of the population. It is seen clinically as reticular, papular, plaque-like, erosive, atrophic or bullous types.¹⁻³

Intraorally, the buccal mucosa, tongue and the gingiva are commonly involved although other sites may be rarely affected. Oral mucosal lesions present alone or with concomitant skin lesions. The skin lesions present as violaceous flat-topped papules in ankles, wrist, and genitalia, but characteristically the facial skin is spared.⁴⁻⁶

Vesicle and bulla are seen in the oral lesions of lichen planus. The disease manifests in the oral cavity several weeks before the skin lesions. About 15% of oral lichen planus patients have concurrent skin lesions. The reticular form has a better prognosis as 40% of cases have spontaneous remission, the erosive type being long standing and with frequent exacerbations and severe pain and complications.⁷

Hence; we conducted the present study to assess the prevalence of OLP in a known population of Samba region, Jammu.

MATERIALS AND METHODS

The present study was conducted at department of skin and V.D. & department of dentistry, S.P. Medical College, Bikaner (Rajasthan) and included evaluation of prevalence rate of OLP. A total of 200 subjects were included in the present study.

Exclusion Criteria

- Patients with history of any systemic illness,
- Patients with any known drug allergy,
- Patients with history of any metabolic disorder,

Detailed demographic details and clinical details of all the patients were recorded. Mouth mirror and probe were used for making the clinical examination of the patients. All the results were recorded on excel sheet. Data were analyzed by SPSS software. Univariate regression curve were used for assessment of level of significance. P-Value of less than 0.05 was taken as significant.

RESULTS

Out of total 200 subjects, 120 were males and remaining were females. Mean age of the subjects of the present study was 38.5 years. OLP was found to be present in 2 males and 1 female. Prevalence rate of OLP was found to be 1.5 percent.

Table 1: Demographic Detail of Patients

Parameter	Values
Mean Age(Years)	38.5
Male	120
Females	80

Table 2: Prevalence Rate of OLP

Prevalence Rate	Values
Male	1.67
Female	1.25
Total	1.5

DISCUSSION

In the present study, we observed that prevalence rate of OLP was 1.5 percent. Varghese SS et al. determined the epidemiology of OLP in a cohort of South Indian population. All the case data records of 29,606 patients who visited Mar Baselios Dental College and Hospital, Kerala, India from 2014 to 2015 were retrospectively reviewed. For data review, 122 patients of OLP were selected. Estimated were type, number, and location of lesions, clinical manifestation, age of the patient, gender, onset and duration of lesion, stressful life style, habits, skin involvement and associated systemic illness, and presence/absence of dysplasia. When the distribution of OLP among the gender was considered, they found more prevalence in females than males. Fifty-seven percent of patients were associated with stressful lifestyle. Reticular lichen planus was the most common clinical subtype found. Bilateral buccal mucosal was the common site, when the distribution of sites of OLP were compared ($P < 0.05$). Anaplasia was found among 5% of lichen planus lesions. OLP patients had high incidence of hypersensitivity reactions and 5% of OLP lesions showed anaplasia.⁸ Omal P et al. evaluated the prevalence of oral, skin, and oral and skin lesions of LP from a population of patients attending the Department of Oral Medicine and Radiodiagnosis, Pushpagiri College of Dental Sciences, Tiruvalla, Kerala, India. A cross-sectional study was designed to evaluate the prevalence of oral, skin, and oral and skin lesions of LP. This is an ongoing prospective study with results of 2 years being reported. LP was diagnosed on the basis of clinical presentation and histopathological analysis of mucosal and skin biopsy done for all patients suspected of having LP. Out of 18,306 patients screened, 8,040 were males and 10,266 females.⁹ LP was seen in 118 cases (0.64%). Increased prevalence of LP was observed in middle age adults (40-60 years age group) with lowest age of 12 years and highest age of 65 years. No statistically significant differences were observed between the genders in skin LP group ($P = 0.12$) and in oral and skin LP groups ($P = 0.06$); however, a strong female predilection was seen in oral LP group ($P = 0.000036$). The prevalence of cutaneous LP in oral LP patients was 0.06%. This study showed an increased prevalence of oral LP than skin LP, and oral and skin LP with a female predominance.¹⁰

Mostafa B et al. reported the prevalence of oral lichen planus among a sample of the Egyptian population. 4470 Egyptian patients, aged 15-75 years, were seen at the outpatient clinic at the Faculty of Oral and Dental Medicine, Cairo University, Egypt. 31.25% of these patients were males and 68.75% were females. Oral mucosal lesions consistent with oral lichen planus (OLP) were identified both clinically and confirmed histologically

(in atypical cases) so that the prevalence of oral lichen planus in this study is 1.43%. 64 patients were diagnosed with OLP (20 males and 44 females). The average age of the affected group was 48.07 years. Associated skin lesions were detected in 15/64 patients (23.44%) and tobacco habits was observed in 20/64 patients (31.25%). The average period of follow-up of the affected cases was 1-2 years, during which two cases developed squamous cell carcinoma of the oral lesion. Within the limitations of this study it revealed the prevalence of OLP among middle-aged women. Atrophic lesions were most frequent, followed by the erosive forms. Anti-HCV circulating antibodies were more common in patients with OLP and, notably, OLP was associated with Diabetes mellitus in 15.63% of patients.¹¹

CONCLUSION

From the above results, the authors concluded that OLP occurs in a significant proportion of population.

REFERENCES

1. Rojo-Moreno JL, Bagan JV, Rojo-Moreno J, Donat JS, Milian MA, Jimenez Y. Psychologic factors and oral lichen planus. A psychometric evaluation of 100 cases. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 1998; 86:687-91.
2. Ismail SB, Kumar SK, Zain RB. Oral lichen planus and lichenoid reactions: Etiopathogenesis, diagnosis, management and malignant transformation. *J Oral Sci*. 2007; 49:89-106.
3. McCartan BE. Psychological factors associated with oral lichen planus. *J Oral Pathol Med*. 1995; 24:273-5.
4. Cutler TP. Lichen planus caused by pyrimethamine. *Clin Exp Dermatol*. 1980; 5:253-6.
5. Ivanovski K, Nakova M, Warburton G, Pesevska S, Filipovska A, Nares S, et al. Psychological profile in oral lichen planus. *J Clin Periodontol*. 2005; 32:1034-40.
6. Hawk JL. Lichenoid drug eruption induced by propranolol. *Clin Exp Dermatol*. 1980; 5:93-6.
7. Hamburger J, Potts AJ. Non-steroidal anti-inflammatory drugs and oral lichenoid reactions. *Br Med J (Clin Res Ed)*. 1983; 287:1258.
8. Varghese SS, George GB, Sarojini SB. Epidemiology of Oral Lichen Planus in a Cohort of South Indian Population: A Retrospective Study. *Journal of Cancer Prevention*. 2016; 21(1):55-59. doi:10.15430/JCP.2016.21.1.55.
9. Omal P, Jacob V, Prathap A, Thomas NG. Prevalence of Oral, Skin, and Oral and Skin Lesions of Lichen Planus in Patients Visiting a Dental School in Southern India. *Indian Journal of Dermatology*. 2012; 57(2):107-109. doi:10.4103/0019-5154.94276.
10. Mostafa B, Ahmed E. Prevalence of oral lichen planus among a sample of the Egyptian population. *Journal of Clinical and Experimental Dentistry*. 2015; 7(1):e7-e12. doi:10.4317/jced.51875.

Source of Support: Nil.

Conflict of Interest: None Declared.

Copyright: © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882. This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article as: Prasoon Soni, Ekta Soni, Jitendra Acharya. A Study of Prevalence of Oral Lichen Planus in a Known Population: A Cross-Sectional Hospital Based Study. *Int J Med Res Prof*. 2018 Nov; 4(6):267-68. DOI:10.21276/ijmrp.2018.4.6.059