

A Study of Thyroid Disorders in Third Trimester of Pregnancy

Mahesh Kumar Goyal¹, Gouri Bhaduri^{2*}, Maheshwar Prasad³

¹Assistant Professor, Department of Preventive and Social Medicine, MGM Medical College, Jamshedpur, Jharkhand, India.

^{2*}Assistant Professor, Department of Physiology, MGM Medical College, Jamshedpur, Jharkhand, India.

³Civil Surgeon- cum - Chief Medical Officer, East Singhbhum Jamshedpur, Jharkhand, India.

ABSTRACT

Background: Thyroid disease is the second most common endocrine disease to affect women of reproductive age. Thyroid disorders can result into adverse reproductive and pregnancy implications. Although gestational hyperthyroidism is uncommon (0.2%), gestational hypothyroidism occurs in higher prevalence (2.5%) and can lead to neonatal and child neurodevelopmental deficits and maternal obstetric complications. Thyroid-related pathophysiologic changes aggravated by pregnancy. Trimester-specific reference intervals for thyroid function tests are critical for maintaining the delicate balance of thyroid hormones during pregnancy. Thyroid-stimulating hormone (TSH) levels are low-normal in the first trimester, with normalization by the second trimester and remains normal in third trimester also. Hypothyroidism during pregnancy is associated with gestational hypertension and low birth weight. Women with hyperthyroidism (high TSH levels) had a >3-fold increase in risk of very preterm delivery.

Objective: To evaluate the prevalence of thyroid disorders in pregnancy.

Methods: This was a observational study among the pregnant women attended the labour room for full term normal delivery. All the pregnant women who were registered at Community Health Centre for delivery along with results of TSH levels were considered as study subjects.

Results: The present study revealed a prevalence of 2.32% of hyperthyroidism and a prevalence of 7.94% of hypothyroidism.

The mean age of study subjects was 23.38 years \pm 3.60. The TSH levels were observed on different parameter like age, caste, parity etc. Most of the pregnant women were primiparous and 2nd gravida.

Conclusions: Thyroid disorders are common in pregnancy, and the most common disorder is subclinical hypothyroidism. Early detection of Thyroid dysfunction can help in starting treatment in affected pregnancies, for better outcome. Hence, Thyroid function test must be advised to all pregnant women.

Key words: Pregnancy, Thyroid Disorders, TSH Levels, Third Trimester.


*Correspondence to:

Dr. Gouri Bhaduri,
Assistant Professor,
Department of Physiology,
MGM Medical College,
Jamshedpur, Jharkhand, India.

Article History:

Received: 29-05-2018, Revised: 21-06-2018, Accepted: 20-07-2018

Access this article online

Website: www.ijmrp.com	Quick Response code 
DOI: 10.21276/ijmrp.2018.4.4.007	

INTRODUCTION

Diseases of the thyroid gland are common; affecting about 5% of the general population, and predominantly affect females. It is a proved fact that maternal thyroid disorder influences the outcome of both mother and fetus, during and also after pregnancy. A normal pregnancy results in a number of important physiological and hormonal changes that alter thyroid function. About 2 to 5% of pregnant woman suffer from any variety of thyroid disorders and timely intervention can be done if detected early.¹ Values of thyroid hormones during pregnancy differ from non-pregnant values because of physiological changes. Values in pregnancy also vary from trimester to trimester, and although no consensus about this value has been made yet. Several community-based cohort studies were conducted to establish trimester-specific reference ranges (TSRRs) for T3, T4, and TSH in different

communities.²⁻⁹ Studies have shown considerable variations in the thyroid hormone profile among populations of different origin, probably owing to modulatory effects of ethnicity¹⁰, parity¹², body mass index (BMI)¹³, iodine insufficiency^{14,15}, and certain pregnancy-induced disorders^{16,17} of thyroid function.

According to recent American Thyroid Association (ATA) guidelines-2017, the recommended reference ranges for TSH are 0.1 to 2.5 mIU/L in the first trimester, 0.2 to 3.0 mIU/L in the second trimester, and 0.3 to 3.0 mIU/L in the third trimester.¹⁸

The most frequent thyroid disorder in pregnancy is maternal hypothyroidism. It is associated with fetal loss, placental abruptions, pre-eclampsia, preterm delivery and reduced intellectual function in the offspring.¹⁹ In pregnancy, overt hypothyroidism is seen in 0.2%²⁰ and sub clinical hypothyroidism

in 2.3% cases.²¹ There has been a wide geographic variation in prevalence of hypothyroidism during pregnancy. It varies from 2.5% from the West to 11% from India.^{22,23} It seems that prevalence of hypothyroidism is more in Asian countries compared to the West.²⁴

Hyperthyroidism occurs in 0.1–0.4% of pregnant women, whereas about 2–3% of pregnant women are hypothyroid, of whom 0.3–0.5% have overt hypothyroidism and 2–2.5% present subclinical hypothyroidism.²⁵ Uncontrolled maternal hyperthyroidism has been associated with fetal tachycardia, small for gestational age babies, prematurity, stillbirths and possibly congenital malformations.

MATERIALS & METHODS

The present observational study was conducted among the pregnant women who came to the Community Health Centre (in municipal area) in third trimester at term for delivery from January 2018 to April 2018. A total number of 763 women were registered for the delivery out of which only 302 were having TSH reports. The data were obtained with regard to age, caste, religion, and parity.

RESULTS

The data was analyzed on SPSS 16.0.

- i) **TSH level:** The American Thyroid Association guidelines 2017 recommends the normal range of TSH in third trimester from 0.3 mIU/L to 3.0 mIU/L. 71.86% of the pregnant women were in the normal range of TSH whereas 2.32% were having a low TSH level (<0.3 mIU/L) i.e. Hyperthyroidism and 25.82% with a high TSH level (>3 mIU/L) i.e. Hypothyroidism (both subclinical and overt). The range of TSH levels varied from a minimum of 0.01 mIU/L to a maximum of 18 mIU/L. However, many of the studies across the globe states that the value of Thyroid varies according to ethnicity. American Thyroid Studies also recommends the use of local data as a reference values. Various Indian studies recommend an upper range of TSH in third trimester of pregnancy at 4.5 mIU/L. However, the American Association of Clinical Endocrinology and the Endocrine Society Consensus panel recommended that 4.5 mIU/L should be maintained as the upper limit of normal. They reasoned that although some individuals within the range of 2.6–4.5 mIU/L may have subclinical thyroid disease, there was a lack of evidence of adverse outcome in this group. Considering this upper value (4.5 mIU/L) the number of pregnant women came to 24 out of 302 pregnant women as Hypothyroidism (7.94%).
- ii) **Age:** The mean age group of the females was 23.38 years \pm 3.60 with a minimum of 18 years and a maximum of 36 years. 58.28% women were in the age group of 19–24 years followed by 31.79% in the age group of 24–29 years. The mean age group of Hyperthyroid pregnant women (n=7) was 23.29 years \pm 5.529 whereas the same among the Hypothyroid pregnant women (n=24) was 22.79 years \pm 3.741.
- iii) **Caste and Religion:** Most of the pregnant women were belonging to the other backward castes (40.7%) and Schedule tribes (37.4%). Muslims were 5.6% among the study subjects. 57.14% were belonging to OBC whereas others were equal 14.28% in Hyperthyroidism category. In

hypothyroid pregnant women, most affected women were 45.33% among OBC and 33.33% among Schedule Tribes.

- iv) **Parity:** Almost 1/3rd (33.1%) of the study subjects were primiparous where as 66.9% were multigravida. Among multigravida most of the pregnant women were either in 2nd gravida (37.4%) followed by 3rd gravida (21.5%). Others were 4th gravida (5.6%) and 5th gravida (2.3%). Most of the hyperthyroid pregnant women (85.17%) were 2nd gravida except one (14.83%) was primiparous whereas in hypothyroid group the group involved was primiparous (29.16%), 2nd gravida (33.34%) and 3rd gravida (29.16%).

DISCUSSION

This present study was aimed to assess the prevalence of thyroid disorders among pregnant women at third trimester. There has been a wide geographic variation in prevalence of hypothyroidism during pregnancy. It varies from 2.5% from the West to 11% from India.

Sahu et al, have done thyroid function during second trimester in high-risk pregnant women and reported that prevalence of thyroid disorders, especially overt and subclinical hypothyroidism was 6.47%.²⁵ In another study from India, Nambiar V et al have reported prevalence of hypothyroidism and thyroid autoimmunity as 4.8% and 12.4%, respectively and were significantly associated with miscarriage²⁷ whereas the present study showed a 2.32% of hyperthyroidism and 7.94% prevalence of hypothyroidism.

CONCLUSION

This study concludes that thyroid disorders are common in pregnancy, and the most common disorder is hypothyroidism. Screening for thyroid function and autoimmunity, and timely and appropriate treatment, will improve pregnancy outcome.

REFERENCES

1. Glinoe D, Soto MF, Bourdoux P, et al. Pregnancy in patients with mild thyroid abnormalities: maternal and neonatal repercussions. *J Clin Endocrinol Metab* 1991;73:421–7.
2. Stricker RT, Echenard M, Eberhart R, Chevaller MC, Perez V, Quinn FA, et al. Evaluation of maternal thyroid function during pregnancy: the importance of using gestational age-specific reference intervals. *Eur J Endocrinol*. 2007; 157:509–14.
3. Soldin OP, Soldin D, Sastoque M. Gestation-specific thyroxine and thyroid stimulating hormone levels in the United States and worldwide. *Ther Drug Monit*. 2007;29(5):553–9.
4. Dhatt GS, Jayasundaram R, Warethc LA, Krishnasamy Jayasundaram NN, Darwish EA, Lewis A. Thyrotrophin and free thyroxine trimester-specific reference intervals in a mixed ethnic pregnant population in the United Arab Emirates. *Clin Chim Acta*. 2006;370:147–51.
5. Azizi F, Mehran L, Amouzegar A, Delshad H, Tohidi M, Askari S, Hedayati M. Establishment of the trimester-specific reference range for free thyroxine index. *Thyroid*. 2013;23(3):354.
6. Marwaha RK, Chopra S, Gopalakrishnan S, Sharma B, Kanwar RS, Sastry A, Singh S. Establishment of reference range for thyroid hormones in normal pregnant Indian women. *BJOG*. 2008;115(5):602–6.
7. Shen FX, Xie ZW, Shi-Ming Lu SM, Tar-Choon Aw TC, Zhu B. Gestational thyroid reference intervals in antibody-negative Chinese women. *Clin Biochem*. 2014;47:673–5.

8. Akarsu S, Akbiyik F, Karaismailoglu E, Dikmen ZG. Gestation specific reference intervals for thyroid function tests in pregnancy. *Clin Chem Lab Med*. 2016. doi:10.1515/cclm-2015-0569. [Epub ahead of print].
9. Bocos-Terraz JP, Izquierdo-Álvarez S, Bancalero-Flores JL, Álvarez-Lahuerta R, Aznar-Sauca A, Real-López E, Ibáñez-Marco R, Bocanegra-García V, Rivera- Sánchez G. Thyroid hormones according to gestational age in pregnant Spanish women. *BMC Res Notes*. 2009;2:237. doi:10.1186/1756-0500-2-237.
10. McLeod DS, Cooper DS, Ladenson PW, Whiteman DC, Jordan SJ. Race/ ethnicity and the prevalence of thyrotoxicosis in young Americans. *Thyroid*. 2015. [Epub ahead of print].
11. Ryckman KK, Spracklen CN, Dagle JM, Murray JC. Maternal factors and complications of preterm birth associated with neonatal thyroid stimulating hormone. *J Pediatr Endocrinol Metab*. 2014;27(9-10):929-38.
12. Solanki A, Bansal S, Jindal S, Saxena V, Shukla US. Relationship of serum thyroid stimulating hormone with body mass index in healthy adults. *Indian J Endocrinol Metab*. 2013;17 Suppl 1:S167-9.
13. Medici M, Korevaar TI, Visser WE, Visser TJ, Peeters RP. Thyroid function in pregnancy: What is normal? *Clin Chem*. 2015;61(5):704-13. doi:10.1373/clinchem.2014.236646.
14. Lazarus JH. Thyroid function in pregnancy. *Br Med Bull*. 2011;97:137-48.
15. Ali AA, Okud A, Khojali A, Adam I. High incidence of obstetric complications in Kassala Hospital, Eastern Sudan. *J Obstet Gynaecol*. 2012;32(2):148-9.
16. Elhaj ET, Adam I, Alim A, Elhassan EM, Lutfi MF. Thyroid function/antibodies in Sudanese patients with preeclampsia. *Front Endocrinol*. 2015;6:87. doi:10.3389/fendo.2015.00087.
17. van den Boogaard E, Vissenberg R, Land JA, van Wely M, van der Post JA, Goddijn M, Bisschop PH. Significance of (sub) clinical thyroid dysfunction a systematic review. *Hum Reprod Update*. 2011;17(5):605
18. Cotzias C, Wong SJ, Taylor E et al. A study to establish gestationspecific reference intervals for thyroid function tests in normal singleton. *Eur J Obstet & Gynecol and Reprod Biol* 2008;137:61-6.
19. Abalovich M, Gutierrez S, Alcaraz G, et al. Overt and subclinical hypothyroidism complicating pregnancy *Thyroid* 2002;12:63-8.
20. Casey BM, Leveno KJ Thyroid disease in pregnancy. *Obstet Gynecol* 2006;108:1283-92.
21. Biondi B, Cooper DS The clinical significance of subclinical thyroid dysfunction. *Endocr Rev* 2008;29:76-13 .
22. Stagnaro-Green A. Thyroid antibodies and miscarriage: Where are we at a generation later? *J Thyroid Res*. 2011; 2011:841949.
23. Männistö T, Väärasmäki M, Pouta A, Hartikainen AL, Ruokonen A, Surcel HM, et al. Perinatal outcome of children born to mothers with thyroid dysfunction or antibodies: A prospective population-based cohort study. *J Clin Endocrinol Metab*. 2009; 94:772-9.
24. Wang W, Teng W, Shan Z, Wang S, Li J, Zhu L, et al. The prevalence of thyroid disorders during early pregnancy in China: The benefits of universal screening in the first trimester of pregnancy. *Eur J Endocrinol*. 2011 Feb; 164:263-8.
25. Negro R, Mestman JH. Thyroid disease in pregnancy. *Best practice & research. Clinical endocrinology & metabolism*. 2011 Dec;25(6):927-43. [PubMed]
26. Sahu MT, Das V, Mittal S, Agarwal A, Sahu M. Overt and subclinical thyroid dysfunction among Indian pregnant women and its effect on maternal and fetal outcome. *Arch Gynecol Obstet* 2010; 281:215-20.
27. Nambiar, Vimal & S Jagtap, Varsha & Sarathi, Vijaya & R Lila, Anurag & Kamalanathan, Sadishkumar & Bandgar, Tushar & Menon, Padmavathy & S Shah, Nalini. Prevalence and Impact of Thyroid Disorders on Maternal Outcome in Asian-Indian Pregnant Women. *Journal of thyroid research*. 2011. 429097. 10.4061/2011/429097.

Source of Support: Nil.

Conflict of Interest: None Declared.

Copyright: © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882.

This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article as: Mahesh Kumar Goyal, Gouri Bhaduri, Maheshwar Prasad. A Study of Thyroid Disorders in Third Trimester of Pregnancy. *Int J Med Res Prof*. 2018 July; 4(4):27-29. DOI:10.21276/ijmrp.2018.4.4.007