

## Study on Impact of Obesity on Lung Volumes in Elderly Population

Dev Kishan Devra<sup>1</sup>, Soniya Pipliwal<sup>2\*</sup>, Bharti Maan<sup>3</sup>,  
Nishanta<sup>4</sup>, Usha Rani<sup>4</sup>, Purnima Rathore<sup>3</sup>

<sup>1</sup>Associate Professor, <sup>2</sup>M.Sc (Medicine) Physiology Final Year,  
<sup>3</sup>M.Sc (Medicine) Physiology, <sup>4</sup>PG Resident (Illrd Year),  
Department of Physiology, S P Medical College, Bikaner, Rajasthan, India.

### ABSTRACT

**Introduction:** The Pulmonary function tests in subjects with normal BMI and with obese individuals were done and correlated with the severity of obesity. Considering the PFT changes, normal study was present in 17%. But, remaining individuals showed proportionate decrease in both FEV1 and FVC values. These changes were independent of age, sex and blood pressure.

**Methods and Results:** Selection of the Subjects Obese individuals of about 30 people with BMI >30 of both sexes of age group between 30 -45 years were included in this study. The healthy volunteers, visitors or relatives of patients visiting S. P. Medical College Hospital, Bikaner Rajasthan were taken as subjects. So, the PFT changes are more of restrictive than obstructive pattern which can leads to decreased chest wall and lung compliance, respiratory muscle strength and endurance which ultimately ends in poor pulmonary ventilation. Hence, this non-invasive procedure may be utilized as a routine screening test for obese people for better medical care.

**Conclusion:** Thus, obesity is associated with a wide variety of PFT abnormalities, many of which can be corrected by weight loss.


**Keywords:** Pulmonary function tests, Obesity, FEV1, FVC.

### \*Correspondence to:

**Soniya Pipliwal,**  
M.Sc (Medicine) Physiology Final Year,  
Department of Physiology,  
S P Medical College, Bikaner, Rajasthan, India.

### Article History:

**Received:** 03-01-2018, **Revised:** 01-02-2018, **Accepted:** 25-02-2018

Access this article online	
Website: <a href="http://www.ijmrp.com">www.ijmrp.com</a>	Quick Response code 
DOI: 10.21276/ijmrp.2019.5.2.047	

### INTRODUCTION

Obesity is a chronic medical condition characterized by excessive accumulation of fat on human body that cause a generalized increase in body mass. Overweight and obesity has become a worldwide health concern. The prevalence of excess weight gain within populations forecasts an increase burden from several diseases, most notably cardiovascular, diabetes mellitus and cancer.<sup>1</sup> The environmental factors such as lifestyle preferences and cultural practices playing pivotal roles in the rising prevalence of obesity, India has now become the third most common country in the world as per the study conducted by Lancet. It is a major risk factor for many acute and chronic disorders, including cardiovascular, cerebrovascular disease and diabetes. It affects the respiratory system as well. It is associated with additional health conditions including carpal tunnel syndrome, deep vein thrombosis and poor wound healing.<sup>2</sup> Obesity has profound effect on physiology of breathing. In brief it can lead to pulmonary compromise in a number of ways including impairment on pulmonary function testing, small airway dysfunction and

expiratory flow limitation, alteration in respiratory mechanism, decreased chest wall and lung compliance, decreased respiratory muscle strength and endurance, decreased pulmonary gas exchange, lower control of breathing and limitation in exercise capacity.

Severe obesity can lead to OHS and sleep apnea with decreased responsiveness to CO<sub>2</sub> and hypoxemia. BMI (Body mass index) is an attempt to quantify the amount of tissue mass in an individual and it is calculated as weight in kilogram divided by square of height in metres.<sup>2</sup>

### MATERIALS AND METHODS

Selection of the Subjects Obese individuals of about 30 people with BMI >30 of both sexes of age group between 30 -45 years were included in this study.

The healthy volunteers, visitors or relatives of patients visiting S. P. Medical College Hospital, Bikaner Rajasthan were taken as subjects and the study was conducted in department of

Physiology, S.P. Medical College Hospital, Bikaner Rajasthan after getting the Ethical committee approval. The subjects were informed about the study and written consent was obtained from them. The selection of obese individual is based on WHO BMI cut off points. Study group consists of obese subjects with BMI >30kg/m<sup>2</sup> without any confounding factors like age, sex and blood pressure and also without any comorbidities like hypertension, diabetes mellitus etc. The control group consists of age and sex matched non obese subjects with BMI of 18.5 to 24.9kg/m<sup>2</sup>. All participants have undergone PFT using spirometry.

#### Study Design

Cross-sectional study

#### Method

Spirometry is an objective, non-invasive and sensitive test which is easy to perform. Before performing the test, the patient was asked to take a deep breath and mouth piece is placed tightly in between the mouth and teeth. Then the patient was asked to completely blow out rapidly into the mouth piece. The parameters like Forced expiratory volume in 1 sec (FEV<sub>1</sub>), Forced vital capacity, Vital capacity, FEV<sub>1</sub>/FEC ratio, Forced expiratory flow were recorded.

#### Statistical Analysis

The data will be subjected to basic statistical analysis (SPSS 2016) and appropriate interpretations will be drawn.

Table 1: BMI

Parameter	Sample size	Mean	Standard deviation	t value	P value
BMI	N1-30	23.133	3.259	10.3773	0.0001***
	N2-26	32.508	3.497		

\*\*\*Extremely statistically significant

Table 2: Parameters recorded

Parameter	Sample size	Mean	Standard Deviation	T value	P value
FVC	N <sub>1</sub> - 30	2.7033	0.6214	2.0082	0.0496**
	N <sub>2</sub> - 26	2.3535	0.6821		
FEV <sub>1</sub>	N <sub>1</sub> - 30	2.2233	0.5112	2.1672	0.0347**
	N <sub>2</sub> - 30	1.9212	0.5309		
FEV <sub>1</sub> /FVC ratio	N <sub>1</sub> - 30	0.81437	0.07350	0.5150	0.6087
	N <sub>2</sub> - 30	0.82462	0.07517		
FEF 25-75%	N <sub>1</sub> - 30	2.4493	0.9738	1.1002	0.2761*
	N <sub>2</sub> - 30	2.1654	0.9508		
PEFR	N <sub>1</sub> - 30	5.0770	1.6737	1.3282	0.1897*
	N <sub>2</sub> - 30	4.4846	1.6539		
FET	N <sub>1</sub> - 30	5.983	1.019	0.1322	0.8953*
	N <sub>2</sub> - 30	6.031	1.579		

\*Statistically not significant

\*\*Statistically significant

## RESULTS AND DISCUSSION

All the obese people included in the present study had not had the PFT changes; some had normal pulmonary function tests (17%). The mean age of obese individuals included in the study was 38.81 years who ranged from 30 to 46 years. But, the earlier studies were carried out only in the older patients (with the mean age of 64 and above who might have had the age-related PFT changes which was not, hither to, ruled out.<sup>3</sup>

Therefore, it could be stated from this study that the obesity is more prevalent in the individuals of middle age groups and the PFT changes noticed in these people are obesity-driven rather than age-related.<sup>4</sup> Our aim of the study is to evaluate PFT abnormalities with varying degrees of obesity. The mean BMI was 33.80 with the range of 30-40. Considering the PFT changes, normal study was present in 17%. But, remaining individuals showed proportionate decrease in both FEV<sub>1</sub> and FVC values.

These changes were independent of age, sex and blood pressure.<sup>5</sup> So, the PFT changes are more of restrictive than obstructive pattern which leads to severe restriction of pulmonary gas exchange. These changes must be considered when evaluating both baseline pulmonary function studies in obese patients and the changes seen during weight reduction. Many of these PFT abnormalities are reversible with substantial weight loss. Thus, obesity is associated with restrictive pattern of pulmonary abnormalities, many of which can be corrected by weight loss.<sup>6</sup>

## CONCLUSION

The influence of obesity causing the restrictive pulmonary dysfunction is proved beyond doubt, from the significant PFT changes. So, this noninvasive procedure may be utilized as a routine screening test for obese people for better medical care. In

addition, the results necessitate further study to evaluate this medical condition thoroughly as a potential risk factor for pulmonary dysfunction.

#### REFERENCES

1. Al Ghobain. The effect of obesity on spirometry tests among healthy non-smoking adults. BMC Pulm Med. 2012 Mar 21;12:10. doi: 10.1186/1471-2466-12-10.
2. Chery MS et al. physiology of obesity and effects on lung function. J Appl Physio.2012,108:206-2011
3. Fulambarkar A et al. Reference values for pulmonary in Asian Indians living in United States. Chest. 2004, 126: 1225-33.
4. Rossiter CE, Weill H. Ethnic differences in lung functions: evidence for proportional differences. IntJ Epidemiol.1974,3:55-61
5. Sin DD, Jones RL, Man SF. Obesity is a risk factor for dyspnea but not for airflow obstruction. Arch Intern Med.2002,162:1477-81.
6. World Health Organization. Obesity: Preventing and managing the Global epidemic 1997, Geneva: WHO.

**Source of Support:** Nil.

**Conflict of Interest:** None Declared.

**Copyright:** © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882.

This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Cite this article as:** Dev Kishan Devra, Soniya Pipliwal, Bharti Maan, Nishanta, Usha Rani, Purnima Rathore. Study on Impact of Obesity on Lung Volumes in Elderly Population. Int J Med Res Prof. 2019 Mar; 5(2):225-27. DOI:10.21276/ijmrp.2019.5.2.047